

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



NOTES: INFORMATION ON THE PCT SYSTEM IS AVAILABLE IN THE PCT HANDBOOK, WHICH IS AVAILABLE FROM THE INTERNATIONAL BUREAU OF PATENT COOPERATION (WIPO), CHAMBERLIN, PENNSYLVANIA, USA.

(43) International Publication Date
20 January 2005 (20.01.2005)

PCT

(10) International Publication Number
WO 2005/005659 A1

(51) International Patent Classification⁷: C12Q 1/68

(21) International Application Number:
PCT/DK2004/000494

(22) International Filing Date: 9 July 2004 (09.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
PA 2003 01070 14 July 2003 (14.07.2003) DK

(71) Applicant (for all designated States except US):
STATENS SERUM INSTITUT [DK/DK]; Artillerivej 5,
DK-2300 Copenhagen S (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PERSSON, Søren
[DK/DK]; Volosvej 7, 1., DK-2300 Copenhagen S (DK).
SCHEUTZ, Flemming [DK/DK]; Sigerstedgade, 1, 2.tv.,
DK-1729 Copenhagen V (DK).

(74) Common Representative: TOFT, Lars; Statens Serum
Institut, Corporate Affairs, Artillerivej 5, DK-2300 Copen-
hagen S (DK).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report
— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: DIAGNOSTICS OF DIARRHEAGENIC *ESCHERICHIA COLI* (DEC) AND *SHIGELLA* SPP.

(57) Abstract: The present invention contains a method for the identification of the diarrheagenic *E. coli* groups: ETEC (entero-
toxicogenic *E. coli*), A/EEC (attaching and effacing *E. coli*) EPEC (enteropathogenic *E. coli*), VTEC (verocytotoxin producing *E. coli*) and EIEC (enteroinvasive *E. coli*), and *Shigella* spp. The bacterial identification is made possible by the specific detection of the following virulence genes: *sta* and *elt* encoding heat stable enterotoxin (ST) and heat labile enterotoxin (LT) characteristic of ETEC, *eae* encoding intimin, characteristic of A/EEC, EPEC or VTEC, *bfpA* encoding bundle forming pilus (BfpA), characteristic of EPEC, *vtx1* and *vtx2* encoding verocytotoxin 1 and 2 (VT1 and 2) characteristic of VTEC, *ipaH* encoding invasive plasmid anti-
gen H (IpaH) characteristic of EIEC and *Shigella* spp., and *ehxA* encoding enterohemolysin (EhxA) characteristic of some EPEC and VTEC strains. The method allows the simultaneous detection of any combination of the 8 virulence genes by one single multi-
plex-PCR. The method is thoroughly validated with respect to sensitivity and specificity, and showed high performance compared to other publication. The method includes an internal positive PCR control and the carry-over prevention system, UNG, which makes it ideal for routine diagnostic analyses. The method can be combined with a number of other technologies leading to even higher sensitivity and reduced time of analysis - both important parameters when diarrheagenic patient or contaminated foods are analysed.